



The Relationship between Sleep Quality, Body Composition, Physical Activity and Depression among Female College Students.

Shokhan Omar Abdulrahman

College of Physical Education and Sports Sciences, University of Sulaimani

shokhan.abdulrahman@univsul.edu.iq

Abstract

There is growing interest in how body composition, physical activity, sleep quality, and depression are interconnected, especially among female college students. This Study aims to enhance current understanding by examining a conceptual model that explores these associations among female college students. This quantitative study involved 132 female college students aged 18 to 24 from the University of Sulaimani (UOS), Kurdistan Region, Iraq. A convenience sampling method was employed, and the public university was selected randomly. Descriptive statistics were computed using SPSS software, and data were analyzed with Partial Least Squares Structural Equation Modeling (PLS-SEM). The results designated an affirmative correlation between Body Mass Index and depression among female college students. Conversely, physical activity exhibited a negative correlation with depression. Additionally, negative correlations were found between sleep duration, sleep efficiency, and depression. Conversely, sleep disturbances and the use of sleep medication were positively correlated with depression. Subjective sleep quality also had a significant negative relationship with depression. However, no significant correlations were observed between body fat, daytime dysfunction, and sleep latency with depression among the participants. This study underscores the complex interplay between Body composition, Physical activity, and sleep quality relative to depression among female college students. Considerably higher depression was associated with elevated BMI and poor sleep-related factors, but better sleep quality and physical activity were linked to lower depression rates. These results point out that interventions aimed at sleep quality, as well as physical training programs, could help minimize depression within this population bracket. However, more research needs to be done to understand these associations better and design precise plans for enhancing mental health.

KEYWORDS: Physical activity, Body composition, Sleep quality, Depression, Female college students.

Received: 27/9/2024

Accepted: 17/10/2024



1- INTRODUCTION

Mental well-being is a fundamental aspect of general health, especially for college students traversing difficult and, consequently, stressful settings. Despite the frequent prevalence of mental state concerns among college students, effective intervention programs are still lacking. It has been shown that mental, emotional, and behavioral disorder rates can go as high as 20% during adolescence life span (Cahuas *et al.*, 2019). Depressive disorder is much more common among all these disorders, making it problematic because it is usually long-term, requires extensive therapy, and has serious effects on recovery. Depressive people in colleges feel insufficient, scared, or indignant all the time (Alam *et al.*, 2023). Reported rates of depression prevalence among college students in India vary significantly, with ranges from 21.5% to 71.25% (Meitei & Singh, 2019). Hence, these changes underscore the importance of targeting counseling services mainly for female students who bear the larger burden. This underscores how important it is to have effective mental health interventions for these people, given that the disease affects their academic performance, social well-being, and general happiness in life (Wilson *et al.*, 2014). Depression can be influenced by various factors like body composition, physical activity, and better SQ. Good sleep can easily prevent depression since it comes with mood disorders (Vanteemar *et al.*, 2019). Body composition (BC), which includes body mass and, to some extent, fat spread, is linked to mental wellness, too, just as research findings indicate that depression can be influenced by being too skinny or overweight (Lv *et al.*, 2024). Depressive symptoms can possibly be reduced through engaging in exercise, which is thought to have mood-elevating effects (Singh *et al.*, 2023). Although there is significant research has been done in investigating how depression is related to the quality of sleep, the composition of the body, and physical activity by separate studies, there is very little corresponding research to show how all these factors are interconnected and contribute to overall depression. Regularly, studies only consider one variable at a time without considering how others might influence it, as well as not looking at their combined effects or possible interactions among them.

Findings indicate that poor sleep quality increases the risk of developing depressive symptoms significantly. People with persistent sleep disturbances are at a higher risk of developing depression (Tanahira *et al.*, 2023). In numerous research, the existence of high amounts of body fat is associated with depression, an aspect that is closely linked to mental health in body composition (Lv *et al.*, 2024; Rindler *et al.*, 2023). Besides, some studies have consistently linked normal physical activity with reduced depression levels, meaning the activity has protection properties. Protection against the development and worsening of this mental health condition through elevation in Physical activity levels has been recorded (Battalio *et al.*, 2020). Depression is multifaceted and has various reasons, making it necessary for a comprehensive technique, which includes SQ, BC, and PA (Viinikainen *et al.*, 2024; Omichi *et al.*, 2022; Guo *et al.*, 2020). The intention of this study model is to disentangle the relationship between these aspects and their impact on depression, with the end in view being to provide information that could help design methodical gender-based interventions that are more effective among female undergraduates. Since female college students are in a rite of passage stage, it is critical to study SQ, body composition, and PA in the depression context of females. The objective of this research is to explore a new model and present a more comprehensive picture of the contributory effects of these factors on mental health that would help in developing better prevention and intervention programs against depression.



2- LITERATURE REVIEW

The Relationship between SQ, BC, and PA and depression among Female college students is not covered soundly in the literature. Especially in the Middle East, this type of research needs to be given greater attention, and inferior studies focused on the association between these factors, especially among female college students. The previous studies were conducted in America, Germany, India, China, and Saudi Arabia. The reviewed studies investigated the interplay between SQ, BC, PA, and depression among college students across various contexts. Previous studies conducted in China (Guo *et al.*, 2020; Zhang *et al.*, 2022; Zhou *et al.*, 2021; Xiang *et al.*, 2020) the studies collectively highlight the connection between PA and mental health outcomes among college students. The study (Guo *et al.*, 2020) demonstrates a link between PA intensity and reductions in depressing symptoms. Furthermore, the study (Zhang *et al.*, 2022) explores how exercise impacts depression through mediators like self-concept and social support. A study (Zhou *et al.*, 2021) focuses on how sedentary behavior and PA relate to depressive indications in a specific population of Sports College students. Additionally, the research (Xiang *et al.*, 2020) “examines the effects of PA on depression and anxiety in the background of the COVID-19 epidemic. Each study focuses on specific subgroups or contexts, such as sports university students, college students during a pandemic, or students involved in particular projects, which may limit the applicability of their results to the broader college student population. The findings may not generalize to other groups due to differences in demographics, cultural contexts, or unique situational factors. Previous research conducted in China and India (Cahuas *et al.*, 2019; Zhang *et al.*, 2024; Shahbaz *et al.*, 2023) has collectively highlighted the connections between PA, SQ, and depression outcomes among college students. A study conducted in China (Cahuas *et al.*, 2019) investigates how PA and sleep are related to depression in college students. It suggests that both regular PA and quality sleep are associated with lower levels of depression. Another Chinese study (Zhang *et al.*, 2024) focuses on the impact of PA on anxiety, depression, and SQ among college students following the COVID-19 pandemic. It emphasizes that PA can moderate the effects of the pandemic on mental health and SQ. Research executed in India (Alam *et al.*, 2023) examines the frequency of PA and sleep and their associated factors in relation to depression among college students. It finds that both the frequency of PA and SQ are linked to depressive symptoms, highlighting the importance of consistent healthy behaviors. These studies (Cahuas *et al.*, 2019) (Zhang *et al.*, 2024) (Alam *et al.*, 2023) highlight specific aspects or contexts (such as frequency of activity, post-pandemic conditions, or combined effects of PA and sleep), which may limit how well the results apply to other groups of college students or different demographic populations. The findings may not be fully generalized due to these focused contexts and varying methodological approaches. A study that was done in the U.S. looked at how the combination of neighborhood aspects, individual practices, and sleep patterns affected cases of depression in college girls. It identified significant correlations between disrupted sleep and increased depressive symptoms, highlighting social and behavioral influences. However, the study, conducted at a secretive women’s university in the southeast U.S., may have limited generalizability to other settings (Wilson *et al.*, 2014). Furthermore, previous studies (Alqurashi *et al.*, 2022; Zhang *et al.*, 2022; Dudo *et al.*, 2022) have collectively underscored the link between sleep patterns and depression in college students. The study undertaken in Saudi Arabia investigated (Alqurashi *et al.*, 2022) how sleep period and quality affect depression amongst US and faculty. It found that shorter sleep length and poorer SQ are strongly linked to higher depressive symptoms, underscoring the need for better sleep management. Research carried out in China (Zhang *et al.*, 2023) Focused on the long-term effects of healthy sleep patterns on depressive trajectories among college students. It revealed that maintaining consistent, healthy sleep patterns can mitigate depressive symptoms over time, suggesting a preventive approach to mental health. An investigation in Germany (Dudo *et al.*, 2022) examined sleep patterns and depressive symptoms, specifically in medical students, emphasizing



the high prevalence of sleep disturbances and their association with depression. A significant gap across these studies (Alqurashi *et al.*, 2022; Zhang *et al.*, 2022; Dudo *et al.*, 2022) is the limited focus on specific dimensions of sleep, such as daytime dysfunction, sleep efficiency, sleep latency, sleep disturbances, subjective SQ, and use of sleeping medications. These dimensions are crucial for a comprehensive understanding of how different aspects of sleep impact depression. The studies generally address broader sleep characteristics without delving deeply into these nuanced elements, which could provide additional insights into the complex relationship between sleep and depression. Evidence from research in the U.S. and Romania (Torres *et al.*, 2023; Caciulal *et al.*, 2024) reveals a significant relationship between PA, BC, and mental health. An examination conducted in the U.S. (Torres *et al.*, 2023) analyzed correlations between mental health, PA, and BC in American college students following the COVID-19 lockdown. It found significant links between decreased PA and worsening mental health alongside changes in BC. Additionally, an investigation compared mental health, BC, and PA between American and Romanian college students post-lockdown. It highlighted similar patterns in both populations, with lower PA and altered BC correlating with poorer mental health (Caciula *et al.*, 2024). A notable gap in these studies is that the first study's focus on American college students limits its applicability to other cultural and regional contexts (Torres *et al.*, 2023). While the second study offers a comparative perspective between American and Romanian students, it must thoroughly address SQ. This omission is significant because SQ can interact with BC and PA to influence mental health (Caciula *et al.*, 2024). Previous studies came from American, German, Romanian, Indian, Chinese, and Saudi Arabian contexts. However, none have been conducted in the Iraqi framework. Owing to the research location, chosen research approach, applied sampling techniques, and the population sample, the earlier findings may not apply in the Iraqi context. However, only one of the previous studies was conducted in the Middle East (Alqurashi *et al.*, 2022). None of the previous studies have explored the model of the relationship between BC, SQ, PA, and Depression among female college students. Additionally, this limitation highlights the issue of restricted generalizability and underscores the need for research specifically targeting female college students within the Iraqi context.

2.1 THEORETICAL MODEL

The Biopsychosocial model: Created in the 1970s by Dr. George Engel and Dr. John Romano (Engel, 1977), this approach encompasses an integrated view of health by amalgamating biological, psychological, and social dimensions. It shows how body composition affects neurotransmitter levels like serotonin and dopamine, which are essential for mood management. Therefore, any unusual fat circulation in our bodies may lead to these chemicals, possibly causing depression. Body image concerns can have subsequent consequences on mood and behavior psychologically. On a social scale, however, emphasis on what a healthy human body should look like may result in certain stresses that can, in turn, aggravate mental illness, such as depression. This design incorporates how physical and psychological elements have affected other parts of health (Merino *et al.*, 2024). From a biopsychosocial perspective, PA exercise has been positively linked to physical, psychological, and social treatments for depression. PA can elevate neurotransmitter control, release endorphins, and reduce inflammation, all of which improve mood and lessen depressive symptoms. On a psychological level, it increases self-esteem, enhances cognitive skills, and makes stress easier to control. From a social perspective, it enhances positive attitudes among people while making them stick to a daily schedule that promotes general mental health. The thorough approach helps in highlighting the various ways through which physical inactivity may enhance depression (Eime *et al.*, 2013). The Sleep-Affect Model (Sleep-Wake Model of Depression) effectively



addresses and confirms the link between SQ and depression. For instance, this model provides an extensive framework through which one can decipher how disturbance in sleep quality worsens depressive symptoms' beginning or duration (Otte, 2009; Riemann *et al.*, 2020; Luo *et al.*, 2024). How do Depression and Poor Sleep Quality (SQ) Interact According to the Model? For example, it has taken into consideration how lower levels of SQ can diminish the ability to handle emotions, hence increasing sensitivity, stress as well and mood instabilities (leading to depression). In contrast, depression often causes sleep problems, such as difficulty falling asleep or waking up too early in the morning. These exacerbate the symptoms, making it even harder for them to break out from themselves. In managing or even preventing depressive disorders successfully then, addressing both slumber times and attitudes becomes crucial (Luo *et al.*, 2024; Luo & Lin, 2024)

3- HYPOTHESES OF THE STUDY

Previous studies support this hypothesis, claiming an existing relation among these variables. This study seeks to establish whether this correlation exists based on observed data or if it delves deeper into divergent perspectives using these connections.

3.1 Body composition

According to past research, BMI and depression have been found to be positively correlated. Negative self-perceptions are among the many negative symptoms that are likely to worsen depressiveness associated with raised BMI. Previous research has shown that weight status may have adverse mental health outcomes, including the likelihood of depression. It has been shown in many studies that a high BMI may lead to higher levels of depressive symptoms while also increasing the likelihood that an individual will have negative self-perceptions, which further increases the incidence rate (Viinikainen *et al.*, 2024; Lv *et al.*, 2024)

H1: Depression has a positive association with BMI among FCS.

Recent research emphasized that young women often suffer from depression and low self-esteem due to high body fat percentage (Akinyemi *et al.*, 2022). Analyzing how body fatness (BMI- BF) affects mental health becomes vital because it provides insights into possible reasons behind association measures. Thus, increased depression levels may be related to greater amounts of body fats, which can, in turn, be attributed to underlying factors like dissatisfaction with appearance, comparing oneself with others socially, or feeling ashamed in society. In society and culture, female college students are surrounded by pressures that are related to the image of the body. As a result, these individuals often feel bad about themselves when they see an idealized version of what is beautiful in their society or on screen: this can lead to low self-esteem or dissatisfaction with one's appearance. Recognizing why someone becomes depressed because she has gained so much weight would make it easier to understand the difficulties faced by young people who struggle with eating disorders (Lv *et al.*, 2024).



H2: Depression is positively associated with BF among FCS.

3.2 Physical activity

According to (Guo *et al.*, 2020; Mu *et al.*, 2024), it has been shown in past research that students who have high overall PA levels are less likely to have symptoms of depression. As an illustration, the findings suggest that people who engage in regular exercises tend to exhibit fewer depressive signs while experiencing less psychoticism as well as positive affective responses than those who do not. Suppose it can comprehend the connection between physical activity and depression. In that case, it will be possible to create some policies that would help enhance mental health and prevent occurrences of depressive disorders. It is possible to reduce the burden of depression by promoting physical activity (PA) and using it as a preventive or curative medicine, thus improving the standard of living of individuals. Female University students need to understand the effects of PA on depression in order for them to come up with gender-specific programs dealing with mental health effectively. Understanding the connection might help define the PA as a coping strategy for depression. At a college level, women can have personalized services like mental health care and fitness regimes that are combined with or only involve PA.

H3: Among female college students, PA is negatively associated with depression.

Sleep Quality

The earlier investigation has discovered a correlation between extending the time for falling asleep and enhanced levels of depression signs (Omichi *et al.*, 2022). This implies that individuals taking more extended periods of sleep might be in more dangerous depressive states. Prolonged onset of sleep can impair mood and thus make the body vulnerable to depression by causing disruptions in SQ. Difficulties in managing academic and social issues due to these impairments can also increase the chances of developing depressive symptoms and, in turn, increase the risk of having depression symptoms at the same time (Griggs *et al.*, 2020). As a result, an increase in sleep latency tends to be significantly associated with elevated levels of depression among females undergoing college education.

H4: Among FCS, depression is positively associated with sleep latency.

Prior research has shown that among university students, reduced sleep duration was linked to higher depression prevalence rates (Li *et al.*, 2020; Zhang *et al.*, 2023). When sleep hours are reduced, one can find oneself facing more stress while also having a reduced ability to handle everyday challenges, especially for ladies in college, where both academics and social life create enormous pressures for most of them. Disordered patterns or deficits in sleep are linked to bad moods, poor emotional control, and, as such, increasing chances of being symptomatic of a depressive disorder (Riemann *et al.*, 2020). Because of this fact, it can be inferred that shorter sleep durations are pertinent factors to the prevalence of depression among female university students.

H5: Amongst FCS, sleep duration is negatively associated with depression.



Earlier studies have indicated that one of the important indicators of the severity of depression is sleep efficiency. According to (Lim *et al.*, 2022; Yan *et al.*, 2022), extensive variations in sleep efficiency across different days attract increased depression episodes. Typically, low sleep efficiency corresponds to fragmented or low-quality sleep that hinders restorative stages of sleep. It then follows that when this happens, mood regulation and emotional stabilization could be interfered with, leading to predisposing factors to depression. Moreover, ineffective management of daily activities is hindered by poor academic performance caused by poor attention from lack of quality night rest and lots of daytime napping, which reduce productivity levels (El Hangouche *et al.*, 2018). As a result, the above reasons may lead to a rise in the amount of depression experienced by women students in college.

H6: There is a negative correlation between depression and sleep efficiency in female college students.

Several investigations have repeatedly documented a strong positive link between depression and sleep disturbance (Barsha & Hossain, 2020; M Salih *et al.*, 2023). When people do not sleep long enough or experience recurrent awakenings during the night, they only manage to have bits of fragmented res, which culminates into sufficient sleeping moments that are necessary for emotional and mental stability. These can affect one's ability to think clearly by making it harder for them to be in any single moment or both feelings' at once (Benkirane *et al.*, 2022). Academic stress may, therefore, be worsened in college-going women, hence reducing problem-solving abilities and leading to increased chances of getting depressed.

H7: There is a positive relationship between depressive disorders and sleep disturbance in FCS.

In 2024, Chery and other researchers looked at why depressed patients use various types of sleeping pills but cannot find sleep (Chery *et al.*, 2024). Drowsiness, emotional changes, and cognitive impairment are a few adverse effects that sleeping medications might cause, which can be devastating to overall mental health and may lead to depression (Li *et al.*, 2014). Moreover, excessive use or dependence makes it difficult for one to feel like doing anything since there is no hope left except taking another pill after some time, which will knock you out into unconsciousness, but only after experiencing a small high first, like cocaine followed by deep sleep; therefore, exacerbating mood disturbances and increasing the risk for depression.

H8: Among FCS, sleep medication is positively associated with depression.

As argued by Shim *et al.* (2019), depression levels are higher in cases where daytime dysfunction is prominent. Impaired concentration fatigue and reduced productivity are some of the signs of daytime dysfunction, which are mainly attributed to low SQ or inadequate rest. These challenges may lead to increased stress levels or feelings of anxiety about life, resulting in depressive symptoms. On another note, reduced daytime functionality can have a negative impact on education performance and relationships with other people, making one feel inadequate and leading to depression (Shim *et al.*, 2019). Additionally, trouble during the day can disrupt normal routines and coping strategies and create a loop of impaired daily life performance that intensifies bad moods while worsening daytime functioning with depressive symptoms.

H9: Among FCS, daytime dysfunction is positively associated with depression.

Past studies have illustrated an alarming relationship between self-rated subjective sleep quality (SQ) and depressive symptoms, in agreement with the earlier ones (Huang *et al.*, 2016; Gerri *et al.* L, 2024). Depressive symptomatology was one criterion used for lower subjective SQ. When people believe that their social status



is below average, it influences their moods and makes them feel sad. Low social status perception can cause attention deficiency, reduce energy levels, and impair physical health (O'Leary *et al.*, 2021).

H10: Subjective SQ is negatively associated with depression among FCS.

Methods and Material

4.1 Procedure

The Ethics Committee of the College of Sport Science and Physical Education/University of Sulaimani approved this study, and each student could decide on whether or not they wanted to take part based on their own interests without any promise of personal or school advantages. Moreover, participants received written instructions concerning the study process as well as their rights before entering into it; hence, they were free to drop out whenever they wanted to. Owing to the anonymity adopted during data collection exercises and the retrospective characteristic of the same, there was no need to apply informed consent for the data collection procedure, as the researcher is a lecturer at the same university. Thus, the researcher was allowed to collect data without procuring informed consent from the University of Sulaimani. This work plan was guided by Helsinki's declaration made in 1964 and last reviewed in Fortaleza, Brazil, in 2013.

4.2 Participants

The G*Power software, which was used to figure out the overall sample size required for the research model, provided the basis for the participants in this study. It is an approach frequently employed in scientific research for calculating the sample size required for the number of hypotheses or predictors. Multiple functions and settings are available in this software that covers nearly every type of analysis consideration (Kang, 2021). The sample size for ten predictive hypotheses, based on G*Power software, was 74. Hence, in this study, the participants comprised 132 full-time female undergraduate students from the 1st to the 4th year, with ages ranging from 18 up to 24 years inclusively, with a mean age of 20.22 (SD = 1.51). Moreover, the study was conducted at the University of Sulaimani (UOS) in the Kurdistan region of Iraq during the second semester, from April 11 to May 2, 2023. To ensure the validity of the assessment instruments and the cognitive capacity of the participants, measurements of PA, SQ, and depression were administered to all students. Additionally, body composition metrics, including BMI and BF, were assessed for all participants. Data were included only from students who provided complete responses. Incomplete responses or those lacking essential data were excluded from the analysis. Participants were randomly selected from various faculties, excluding those from the department. This separation was due to the unique physical exercise classes undertaken by students in the College of Sport Science and Physical Education, which could cause differences in physical fitness compared to students in other faculties.

4.3 Instruments

4.3.1 Body composition

To determine body fat and BMI, the researcher measured body fat (%) and body mass index (BMI). A body



composition analyzer with height-weight capabilities consisting of an electronic body fat scale (EW-700A) was used to determine body fat percentage (%), BMI, weight (kg), and height (cm). This instrument can precisely measure these parameters. Throughout the process of measurement, participants were told to wear light clothes, take off any metal objects that touched them, keep their backs straight, and avoid talking or changing their positions. The BMI and body fat percentage are variables of interest in this study. The average BMI for participants was 23.15 (SD = 3.72), and the average body fat percentage was 20.32% (SD = 5.22).

4.3.2 Physical activity

The IPAQ-S7S was used to assess PA levels. The Cronbach's alpha coefficient was 0.70. The validity and reliability of IPAQ-S7S have been tested among university students (Craig *et al.*, 2003; Dinger *et al.*, 2006). Thus, the IPAQ-S7S measures the amount of time spent engaging in physical activity during the preceding week by gauging the frequency and duration of staying active through walking, moderate-intensity as well as vigorous-intensity exercises lasting at least ten minutes each. Using the IPAQ data processing guidelines, we estimate individual average levels of PA, which are provided in detail in another place (Craig *et al.*, 2003). Only cases that were invalid in all three PA domains were removed using the case removal approach. Compute total weekly time spent walking and in activity with moderate- and/or high-intensity levels as follows: Multiply days per week per category by time spent daily on days when the activity took place. The amount of time each day devoted to an area is multiplied by the metabolic equivalents of tasks specific to the area, such as walking (3.3 METs), moderate (4 METs), and vigorous (8 METs) exercise and quantified into metabolic equivalents (METs) thus obtaining the physical activity Economic Commission for Africa (ECA) countries per week. A survey of the population showed that there were three levels of PA: low (meaning an individual does not do much exercise), moderate, and high. These categories entail what the established scoring criteria are; they are obtained from the <http://www.ipaq.ki.se> website, and normally for adult (18-65 years) populations involve moderately intense activity for 30 minutes or more for at least 5 days a week or vigorous activity for 20 minutes or more at least three days a week. Categorization: low: does not meet minimum criteria for either moderate or high. Moderate: covers any of the following three concepts: (a) 3 days of at least 20 minutes of vigorous physical activity per day; (b) walking moderately for more than 30 minutes each day on 5 days each week or for each time; or (c) any combination of walking, moderate or vigorous physical activity for 5 days each week with an accumulation of >600 MET-minutes/week. High: meets either of the following two conditions: (a) engaging in high-intensity weekly for >3 days and accumulating ≥ 1500 MET-minutes/week; or (b) 5 days of any combination of walking, moderate-intensity or vigorous-intensity physical activities achieving at least 3000 MET-minutes/week.

4.3.3 Sleep Quality

The most commonly used scale in medical and non-medical populations is the PSQI to assess SQ. It has a Cronbach's alpha value of around 0.83, signaling high reliability. The PSQI has been found to be reliable and valid across various populations (Mollayeva *et al.*, 2016; Wolf & Rosenstock, 2017). Furthermore, it has been shown that even with small sample sizes, the PSQI is valid (Mollayeva *et al.*, 2016). The PSQI is a self-rated questionnaire comprised of nineteen items aimed at assessing SQ for the month preceding it (Buysse *et al.*,



1989). The PSQI includes various aspects of SQ, which are subjective SQ, sleep latency, sleep duration, sleep efficiency, sleep disturbance, use of sleep medication, and daytime dysfunction. These domains range from 0 to 3. After summing up personal scores to obtain a global score, this ranges from 0 to 21, with higher global scores indicating worse SQ.

4.4.4 Depression

In this study, the Zung Self-Rating Depression Scale (SDS) was used as a norm-referenced measure for screening adults for depressive disorders (Zung *et al.*, 1965). It has a Cronbach's alpha value of around 0.79, which is acceptable. That Lee and colleagues utilized it in the study of several populations has been demonstrated clearly by available data for such use in research done previously by Chinese scholars (Lee *et al.*, 1994; Li *et al.*, 2020), also suitable for undergraduates (Nie *et al.*, 2017; Shao *et al.*, 2020). The SDS is a standard instrument intended for self-administration that is meant to gauge the severity of someone's depressive symptoms. The first scale has 20 questions that measure depressive symptoms around these four main factors: anxiety, cognitive-affective (emotional), somatic-vegetative (biological or physiological), and core-depressive symptoms (Kirkby *et al.*, 2005; Dunstan & Scott, 2019). The study adjusted the SDS culturally to conform to the context of Iraqi society. One item about sexual pleasure was omitted because it was considered sensitive and likely to conflict with Iraqi culture. Thus, this modification aimed to enhance cultural understanding and boost both acceptance and precision of depression assessment among the Iraqis. The SDS has been altered to include nineteen items. Ten of them assess negative experiences, e.g., "I feel down-hearted and blue," while the remaining nine determine positive experiences and are scored differently; for example, "I eat as much as I used to eat." Every item was rated by respondents according to their experiences throughout the previous week on the Likert scale. Response options for the negative posers will range from 1 (some of the time) through 4 (most of the time). In the positive items, the frequency of responses varies, going from 4 (some of the time) to 1 (most of the time). The total score is constituted by all 19 items, thereby showing a person's level of depression. Each participant's score is between 25 and 100, where high scores indicate more symptoms. Those with 25 – 49 have average results where else; anything at 70 upwards is generally an indication that one has a terrible case of depression.

4.4 Data collection

The language translation was done using a standard back-translation approach (Cha *et al.*, 2007; Klotz *et al.*, 2023). Physical activity, Sleep quality, and depression were assessed through a systematic translation methodology. The translating procedure required several significant phases. Based on (Carlson, 2000), the translator should be fluent in both languages and informed on the instrument objective. However, this procedure necessitated the involvement of three translators. A multilingual expert utilized the forward translation method to convert the scale from English to Kurdish. A further bilingual expert translated the English text into Kurdish as part of a forward translation process to ensure consistency and accuracy. Content validity has also been applied, an expert evaluation technique to guarantee the validity of the instrument and ensure the translation correctly presents the original concept. The multi-step process guarantees that the translation tool is perceived as reliable and genuine by the intended audiences.



STATISTICAL ANALYSIS

The primary step in the process of data analysis was writing the variables and questions in the code. Next, the data cleaning and preparation involved checking for outliers, suspicious response patterns, missing data, and normalcy using the Statistical Package for the Social Sciences (SPSS) software version 29. Once the data was cleaned, the sample's central tendencies (mean and standard deviations for continuous variables) and the frequency analysis (for categorical variables) were calculated. Using SmartPLS Software version 4.1, the researchers used the SEM approach to measure the premises and model variables. The measurement validity may be evaluated using 0.6 for the model as well as the path model as long as it is in line with the research objectives of explaining and predicting the difference between constructs. SmartPLS is a powerful tool for assessing model prediction hypotheses and models due to its advanced PLS-SEM functionality. Among the features is an intuitive user interface, ample model assessment tools, PLSpredict to assess predictive performance, and efficient data processing. The essence of these traits is to ensure that it is ideal for accurate predictions and complex models since they are both important in disciplines as diverse as computer science and digital marketing (Cheah *et al.*, 2024).

RESULTS

Age and body composition

The data analytics on the female undergraduates indicated that the ages of the students ranged between 18 and 24 years at an average of 20.22 years old (Standard Deviation = 1.51). The Body Mass Index (BMI) ranges from 14.10 to 32.10, with an average of (M=23.15, SD=3.72). The range of body fat percentage (BF) is from 10.30 to 36.50, with an average of (M=20.32, SD= 5.22).

Table 1 Descriptive analysis for age and body composition

	Minimum	Maximum	Mean	Std. Deviation
AGE	18	24.00	20.22	1.51
BMI	14.10	32.10	23.15	3.72
BF	10.30	36.50	20.32	5.22

The results for students' PA showed that 32.6% of students had low PA, 47.7% of students participated in moderate PA, and only 19.7% had high PA (Figure 1).





Figure 1. Distribution of physical activity among students

Sleep Quality

SQ contains seven domains (sleep latency, sleep efficiency, sleep duration, SQ, sleep disturbance, daytime duration, and use of medicine). The first four domains were assessed using different scales, while the last three domains were assessed using similar scales. The results showed that for the sleep latency domain, the highest percentage belonged to 16-30 minutes (36.4%), while 31.8% of students experienced a sleep latency of over 60 minutes. Results for the second domain, sleep efficiency, indicated that it was distributed predominantly in the range of 75-84% (47%) and only 18.9%, reaching over 85% efficiency. Findings for sleep duration showed that most of the students had a sleep duration between 5-6 hours (29.5%), while 20.5% slept over 7 hours. For SQ, results revealed that 30.3% of students responded to it as fairly good and 25% as very good; however, 28% rated their SQ as very bad.





C: Sleep duration	D: Sleep quality
-------------------	------------------

Figure 2. Distribution of students' response to sleep latency, sleep efficiency, sleep duration and sleep quality

The findings for sleep disturbances (table 2), which includes 9 indicators, show that a considerable portion of students experienced sleep disturbances regularly. For instance, 31.1% of the students wake up between three or more times every week in the middle of the night or early in the morning. Results revealed that 32.6% of students had bad dreams. Using a lower perplexity and higher burstiness, it was also a common problem to get up for the bathroom for 55.3% of students. This was something that happened less than once per week, according to the survey results. Other sleep disturbances, such as feeling too cold or coughing/snoring loudly, are less frequent, with 13.6% and 37.1%, respectively.

According to these results (table 2), it was found that for daytime dysfunction, 58.3% of students reported that in no time at all, they found it easy to keep awake while driving, meal taking, or socializing. However, keeping enthusiasm is a challenge for 29.5% of students, which indicates they struggle with these three or more times per week. Even though these challenges, a considerable portion of students (25%) experienced over a one-week period, less than once did 46.3% face problems staying focused, while 28.8% found it hard to stay excited and energized. For using sleeping medication, results showed that it was relatively low among the students. For the majority of students (63.6%), it was reported that no sleeping medication was used in the past month.

Table 2 Sleep disturbances and daytime dysfunction were frequently responded to by students, in addition to medication use.

Domain	Item	Not during the past month	Less than once per week	Once or twice per week	Three or more times per week
Sleep disturbances	Cannot get to sleep within 30 minutes	14.4	32.6	31.1	22
	Wake up in the middle of the night or early morning	17.4	29.5	22	31.1
	Have to get up to use the bathroom	22.7	55.3	18.2	3.8
	Cannot breathe comfortably	34.1	34.1	17.4	14.4
	Cough or snore loudly	37.1	43.9	15.2	3.8
	Feel too cold	13.6	35.6	24.2	26.5
	Feel too hot	26.5	19.7	25.8	28
	Have bad dreams	5.3	43.2	18.9	32.6
	Have pain	18.2	28	34.8	17.4



Daytime dysfunction	During the past month, how often have you had trouble staying awake while driving, eating meals, or engaging in social activity?	58.3	25	15.2	1.5
	During the past month, how much of a problem has it been for you to keep up enthusiasm to get things done?	12.9	28.8	28.8	29.5
Sleeping medication	Use of sleeping medication	63.6	15.9	16.7	3.8

Depression

The results of descriptive statistics for depression domains (anxiety, cognitive, somatic, and core depressive symptoms) are presented in Table 3. According to these results, the highest mean score belonged to core depressive with a median score of $M= 2.766$, $SD=0.449$, followed by cognitive domain shows a median score of $M=2.485$, $SD=0.478$ plus somatic symptoms with a mean score of $M=2.432$, $SD=0.576$. It was found that the lowest mean score was observed for anxiety, with a median score of $M=2.177$, $SD=0.653$.

Table 3 Results of descriptive analysis for depression scale

Sub-domain	Minimum	Maximum	Mean	SD
Anxiety	1.000	3.670	2.177	0.653
Cognitive	1.250	3.750	2.485	0.478
Somatic	1.330	3.670	2.432	0.576
Core depressive	1.570	3.710	2.766	0.449

Measurement model assessment

First-Order Model

All variables in this research were considered as single items, including (BMI, body fat, PA, and seven domains of SQ). Only depression was a second-order construct with four sub-domains. Therefore, the measurement model was used to assess this construct. The results (Table 4) indicated that all subscales of depression (Anxiety, Cognitive, Core Depressive, and somatic) have a high level of convergent validity. Results showed that all items had an acceptable loading factor (above the threshold of 0.5). The research indicates that both the composite reliability (CR) and alpha Cronbach were above 0.7, and the average variance extracted (AVE) surpassed the acceptable threshold (0.5). The HTMT criterion, established by (Henseler et al., 2015), is crucial in evaluating discriminant validity.



Table 4 Convergent validity and reliability assessment of measurements

Construct	No items	Loading	Cronbach's Alpha	CR	AVE
Anxiety	3	0.791- 0.867	0.772	0.782	0.687
Cognitive	4	0.688-0.772	0.717	0.733	0.537
Core Depressive	7	0.589-0.776	0.802	0.812	0.459
Somatic	3	0.583-0.870	0.638	0.732	0.57

Second order assessment

Since depression was a second-order construct, therefore bootstrapping was used to evaluate the second-order model for this construct. Results (Table 5) show that the four sub-dimensions of depression, including anxiety ($l = 0.637$; $p < 0.001$), cognitive ($l = 0.612$; $p < 0.001$), core depressive ($l = 0.853$; $p < 0.001$) and somatic ($l = 0.520$; $p < 0.001$) significantly boosted depression, working under a second-order hidden parameter.

Table 5 Second-order assessment of measurements using Bootstrap

First order	Loading	SE	T value	P Values
Anxiety	0.637	0.055	11.537	<0.001
Cognitive	0.612	0.077	7.989	<0.001
Core Depressive	0.853	0.03	28.573	<0.001
Somatic	0.52	0.079	6.54	<0.001

Discriminant validity was evaluated using the HTMT method, and results (Table 6) indicated that HTMT values were below 0.85, ranging from 0.011- 0.635. Therefore, the variables exhibit significant discriminant validity.



Table 6 Results of Discriminant Validity using HTMT method

	Anxiety	BMI	Body Fat	Cognitive	Core Depressive	Daytime_dysfunction	Physical activity	Sleep_duration	Sleep_efficiency	Sleep_latency	Sleep_disturbances	Somatic	Subjective_sleep quality
Anxiety													
BMI	0.549												
Body Fat	0.46	0.635											
Cognitive	0.248	0.393	0.3										
Core Depressive	0.456	0.558	0.605	0.447									
Daytime_dysfunction	0.102	0.107	0.118	0.132	0.127								
Physical activity	0.326	0.314	0.346	0.361	0.49	0.123							
Sleep_duration	0.106	0.156	0.235	0.119	0.368	0.351	0.026						
Sleep_efficiency	0.487	0.378	0.263	0.180	0.399	0.176	0.371	0.100					
Sleep_latency	0.12	0.029	0.038	0.26	0.123	0.121	0.166	0.073	0.089				
Sleep_disturbances	0.387	0.378	0.412	0.24	0.401	0.39	0.124	0.339	0.132	0.217			
Somatic	0.401	0.381	0.462	0.393	0.323	0.246	0.198	0.296	0.186	0.502	0.393		
Subjective_sleep quality	0.217	0.198	0.325	0.297	0.452	0.015	0.276	0.202	0.206	0.407	0.066	0.438	
Use of sleeping_medication	0.323	0.359	0.315	0.268	0.349	0.144	0.294	0.265	0.286	0.011	0.019	0.139	0.14




Figure 3 Structural model using bootstrapping methods

Path model:

In order to test the main research hypotheses, the path model was tested using the bootstrap approach with 5000 samples. The collinearity assumption indicates that there were no multicollinearity issues among predictors, and the highest VIF value was 2.136 (below 2.5). The results of the path model using Bootstrap (Table 7) presented that there was a positive and important association between BMI and depression ($\beta = 0.223, P = 0.003, f^2 = 0.085$), indicating that higher BMI is related to advanced stages of depression. There was a negative and significant relationship between PA and depression ($\beta = -0.171, P = 0.006, f^2 = 0.073$). Similarly, the result also showed that there was a negative and significant relationship between sleep period ($\beta = -0.174, P = 0.006, f^2 = 0.069$) and sleep efficiency ($\beta = -0.150, P = 0.019, f^2 = 0.055$) with depression. Results indicate that there were positive and significant relationship between sleep disturbances ($\beta = 0.272, P < 0.001, f^2 = 0.149$) and use of sleeping medication ($\beta = 0.156, P = 0.019, f^2 = 0.058$) and depression while there was a negative and significant relationship between subjective SQ ($\beta = -0.202, P < 0.001, f^2 = 0.090$) plus depression. These results also showed that there was no significant relationship between Body Fat ($\beta = 0.112, P = 0.150, f^2 = 0.020$), Daytime impairment ($\beta = 0.040, P = 0.478, f^2 = 0.004$), and sleep latency ($\beta = -0.051, P = 0.441, f^2 = 0.007$) with depression. The adjusted R square (R2) value depression as a dependent variable was 0.679, revealing that 67.9% SQ, PA, and BMI explained such a huge amount of variance in depression (Figure 3).



Table 7. Path model assessment using Bootstrap.

Path	β	SE	T value	P values	f ²
BMI -> Depression	0.223	0.075	2.976	0.003	0.085
Body Fat -> Depression	0.112	0.078	1.44	0.15	0.02
Daytime_dysfunction -> Depression	0.04	0.057	0.709	0.478	0.004
Physical activity -> Depression	-0.171	0.062	2.774	0.006	0.073
Sleep_duration -> Depression	-0.174	0.063	2.763	0.006	0.069
Sleep_efficiency -> Depression	-0.150	0.064	2.346	0.019	0.055
Sleep_latency -> Depression	-0.051	0.067	0.771	0.441	0.007
Sleep_disturbances -> Depression	0.272	0.074	3.651	0	0.149
Subjective_sleep quality -> Depression	-0.202	0.062	3.248	0.001	0.090
Use of sleeping_medication -> Depression	0.156	0.066	2.354	0.019	0.058

DISCUSSION

The current research indicated a negative and significant relationship between PA and depression. This suggests that higher levels of physical activity among students are strongly connected to a notable decrease in depression rates. It was indicated by previous research that greater overall PA quantities were linked to lesser proportions of ill health symptoms in college students (Zhang *et al.*, 2024; Guo *et al.*, 2020). This is because female college students might have increased satisfaction with their body image; young women could actually end up reaping greater psychological advantages, and well-being when it comes to engaging in physical activities. This result simply shows that physical exercise helps to reduce the severity of depression, raise satisfaction with life, improve mood, strengthen strong then and strengthen the sense of purpose and meaning in life among people. Furthermore, in this present investigation study, sleep duration is negatively correlated with depression. The findings suggest that sleep duration negatively impacts the prevalence rate of depressive symptoms among students, therefore inferring that quality and quantity of sleep are negatively associated with students' mental health. Additionally, other studies also found that among college students, a high proportion are depressed where there is too little sleep (Li *et al.*, 2020; Zhang *et al.*, 2023). In the same vein, depression had an inverse relationship with sleep efficiency (SE). It can be argued that lower depression cases are associated with higher sleep efficiency levels. Additionally, our results concur with other studies that assert that sleep efficiency is a key predictor of depression's severity, while marked fluctuations in daily SE predict higher levels of depression and anxiety among young adults (Lim *et al.*, 2022; Yan *et al.*, 2022). Therefore, it is pertinent that people maintain a high SE to prevent depression symptoms.

In this study, a positive association between sleep disturbances and depression was found. The outcome is in agreement with an earlier study that was conducted by 416 Iraqi adults, which highlighted the strong positive correlation between sleep disturbance and depression symptoms (M Salih *et al.*, 2023). Additionally, factors such as sleep disturbances have been linked to a higher likelihood of depression among young women (Barsha & Hossain, 2020). Moreover, research indicates that women are more disposed to mental health issues like depression (Salk *et al.*, 2017). Collectively, these findings underscore that sleep disturbances and depression among women are influenced by a range of factors, including psychological differences, hormonal fluctuations, and societal pressures. This study found that the Use of sleeping medication positively correlated



with depression among members of this specific sample. Research supports a positive relationship between the use of sleeping medication and depression. Individuals experiencing depression often suffer from poor SQ and are more inclined to use sleep medications (Chery *et al.*, 2024), findings that align with our study results. Thus, this dynamic between sleep medications, depression, and their associated symptoms illustrates the complex interplay between SQ, mental health, and the use of medication.

The participants who participated in this study reported a low subjective SQ that coincided with an increased case of depression; according to the evidence gathered, depressive symptoms are more severe when the quality of sleep is low. Previous research findings (Huang *et al.*, 2016; Gerri *et al.*, 2024; Nakajima *et al.*, 2023) noted that across numerous psychiatric disorders, there is an association between subjective insomnia and depressive symptoms, and this suggests those were two across diagnostic features in different modalities. Also, this study established what women often find challenging in terms of juggling between work, taking care of kids or family members, including house chores; hence, it increases their likelihood of experiencing anxiety attacks coupled with depression, disrupting their sleeping patterns. Increased interventions for better sleep and less depressive symptoms are really needed among diverse ages and social categories to understand the healing process. The current study found no significant association between body fatness (BF %) and depression. The outcome agreement from another study found that body fat distribution was not significantly associated with depression in children and adolescents in the age brackets (Yuan *et al.*, 2023). Together, these findings suggested that the association between depression and body fat may be multifaceted. Despite the suggestion that certain aspects of body fat distribution can affect one's chances of developing depression, its general correlation is hard to define because it differs across different ages or between males and females. There were also no significant associations between daytime dysfunction and sleep latency with depression. Hence, it may be possible that the reason for not finding a relationship between daytime dysfunction and sleep latency with depression in our study was the inadequate sample size. A bigger sample size could uncover linkages that were too small for our research to show. Furthermore, results might be affected by characteristics like age, sex, or certain population features in the sample, which makes them not applicable to other people. For instance, if the sample had a low prevalence of severe daytime dysfunction or depression, this might have affected the results. As for this study, it presents some advantages and disadvantages. The strengths of this study have made significant contributions to understanding the interplay between physical activity, body composition, SQ, and mental health and has introduced a novel exploratory model. While previous research has highlighted these factors individually, this study is unique in that it addresses the specific issues faced by female university students in Iraq, an area that has not been extensively explored. By investigating these challenges, the study provides new longitudinal evidence and explores a conceptual model that maps out the relationships among PA, BC, SQ, and depression in this demographic. In doing so, it addresses a serious gap in the existing literature and advances our awareness in this field. While there are limitations to the research in terms of these issues, a more in-depth survey with 132 participants would enhance its accuracy. This would lead to more reliable results for wider use, considering it would be replicated and improved on in future studies. In addition, it should be noted that respondents came only from one higher education institution situated locally within Iraqi Kurdistan, which makes it difficult for others to accept these findings worldwide. Future research should focus on a variety of demographics over time to understand the connection between these success factors. The implications of the results of this study are essential indicators for academic proceedings and public wellness. By clearing up the association between body composition, sleep quality, physical activity, and depression among female college students, this study highlights the significance of comprehensive health strategies in the educational environment. Educational foundations can take advantage of these viewings to improve aimed interference



that develops better sleep quality, motivates engaging physical activity, and stimulates psychological well-being. Besides, the research's findings could contribute to upcoming studies by determining essential domains for additional inquiry, including the role of sleep treatment in promoting mental wellness results. Decision-makers can also utilize these results to recommend integrated wellness strategies that respond to the multiple issues faced by undergraduate students, eventually helping to develop health and educational achievement.

Conclusion

The relationship between Body composition, Physical Activity, Sleep quality, and Depression among female college students is very complex, according to the research. Higher depression levels were associated with elevated BMI and poor sleep-related factors, as opposed to lower depression being linked with increased PA and better sleep quality. This study, therefore, suggests that there might be a need for intervention programs that focus on improving physical activity levels in order to reduce depression symptoms among female students in their tertiary levels of study. It is important to note that sleep should be considered during treatment for depression, particularly when there is an outbreak of a public health crisis. It is recommended that in order to fully comprehend how daytime dysfunction relates to depression, different variables should be investigated in larger and more varied samples using longitudinal designs. These results provide novel continuous proof for the prompt establishment of a conceptual model representing the correlation between Body composition, physical activity, sleep quality, and depression in female college students.

REFERENCES

- Akinyemi, O.A. et al. (2022). 'Association Between Obesity and Self-Reported Depression Among Female University Students in the United States', *Cureus*, 14(11), pp. 1–9. doi:10.7759/cureus.31386.
- Barsha, R.A.A. and Hossain, M.B. (2020). 'Trouble sleeping and depression among US women aged 20 to 30 years', *Preventing Chronic Disease*, 17, pp. 2009–2016. doi:10.5888/PCD17.190262.
- Battalio, S.L., Huffman, S.E. & Jensen, M.P. (2020). 'Longitudinal associations between physical activity, anxiety, and depression in adults with long-term physical disabilities', *Health Psychology*, 39(6), pp. 529–538. doi:10.1037/hea0000848.
- Benkirane, O. et al. (2022). 'Impact of Sleep Fragmentation on Cognition and Fatigue', *International Journal of Environmental Research and Public Health*, 19(23), pp. 1–20. doi:10.3390/ijerph192315485.
- Buysse, D.J. et al. (1989). 'The Pittsburgh Sleep Quality Index: A New Instrument for Psychiatric Practice and Research', 28(2), pp. 193–212. doi:10.1016/0165-1781(89)90047-4.
- Cahuas, A. et al. (2019). 'Relationship of physical activity and sleep with depression in college students', *Journal of American College Health*, 68(5), pp. 1–8. doi:10.1080/07448481.2019.1583653.
- Carlson, E.D. (2000). 'A Case Study in Translation Methodology Using the Health- Promotion Lifestyle Profile II', 17(1), pp. 61–70.



- Cha, E.S., Kim, K.H. and Erlen, J.A. (2007). 'Translation of scales in cross-cultural research: Issues and techniques', *Journal of Advanced Nursing*, 58(4), pp. 386–395. doi:10.1111/j.1365-2648.2007.04242.x.
- Cheah, J.-H., Magno, F. & Cassia, F. (2024). 'Reviewing the SmartPLS 4 software: the latest features and enhancements'. Springer. doi:10.1057/s41270-023-00266-y.
- Chery, M.J. et al. (2024). 'Depression, Sleep Health & Sociodemographic Correlates in a Nationwide Survey: Implications for Depression Treatment During the COVID-19', *Nature and Science of Sleep*, 16(December 2023), pp. 17–31. doi:10.2147/NSS.S434148.
- Craig, C.L. et al. (2003). 'International physical activity questionnaire: 12-Country reliability and validity', *Medicine and Science in Sports and Exercise*, 35(8), pp. 1381–1395. doi:10.1249/01.MSS.0000078924.61453.FB.
- Dinger, M.K., Behrens, T.K. and Han, J.L. (2006). 'Validity and reliability of the international physical activity questionnaire in college students', *American Journal of Health Education*, 37(6), pp. 337–343. doi:10.1080/19325037.2006.10598924.
- Dudo, K. et al. (2022). 'The association of sleep patterns and depressive symptoms in medical students: a cross-sectional study', *BMC Research Notes*, 15(1), pp. 4–9. doi:10.1186/s13104-022-05975-8.
- Dunstan, D.A. and Scott, N. (2019). 'Clarification of the cut-off score for Zung's self-rating depression scale', *BMC Psychiatry*, 19(1), pp. 1–8. doi:10.1186/s12888-019-2161-0.
- Eime, R.M. *et al.* (2013). 'A systematic review of the psychological and social benefits of participation in sport for adults: Informing development of a conceptual model of health through sport', *International Journal of Behavioral Nutrition and Physical Activity*, 10(1), pp. 1–14. doi:10.1186/1479-5868-10-135.
- Engel, G.L. (1977). 'The Need for a New Medical Model: A Challenge for Biomedicine', *Science*, 196(4286), pp. 129–136. doi:10.1126/science.847460.
- Gerri Satvik Emmanuel and Lokesh L (2024) 'The relationship between sleep quality and depressive symptoms among young adults of India', *World Journal of Advanced Research and Reviews*, 22(1), pp. 487–493. doi:10.30574/wjarr.2024.22.1.1088.
- Guo, F. et al. (2020). 'Intensity of physical activity and depressive symptoms in college students: Fitness improvement tactics in youth (fityou) project', *Psychology Research and Behavior Management*, 13(8), pp. 787–796. doi:10.2147/PRBM.S266511.
- El Hangouche, A.J. *et al.* (2018). 'Relationship between poor quality sleep, excessive daytime sleepiness and low academic performance in medical students', *Advances in Medical Education and Practice*, 9, pp. 631–638. doi:10.2147/AMEPS162350.
- Henseler, J., Ringle, C.M. and Sarstedt, M. (2015). 'A new criterion for assessing discriminant validity in variance-based structural equation modeling', *Journal of the Academy of Marketing Science*, 43(1), pp. 115–135. doi:10.1007/s11747-014-0403-8.
- Huang, V. et al. (2016). 'Subjective sleep quality as a possible mediator in the relationship between personality



- traits and depressive symptoms in middle-aged adults', *PLoS ONE*, 11(6). doi:10.1371/journal.pone.0157238.
- Kang, H. (2021). 'Sample size determination and power analysis using the G * Power software', *Journal of Educational Evaluation for Health Professions*, 18, pp. 1–12. doi:10.3352/jeehp.2021.18.17.
- Kirkby, R., Al Saif, A. & el-din Mohamed, G. (2005). 'Validation of an Arabic translation of the Zung self-rating depression scale', *Annals of Saudi Medicine*, 25(3), pp. 205–208. doi:10.5144/0256-4947.2005.205.
- Klotz, A. C., Swider, B. W., & Kwon, S.H. (2023). 'Back-translation practices in organizational research: Avoiding loss in translation', *Journal of Applied Psychology*, 108(5), p. 699.
- Lee, H.C. et al. (1994). 'The Zung Self-Rating Depression Scale: Screening for Depression Among the Hong Kong Chinese Elderly', *Journal of Geriatric Psychiatry and Neurology*, 7(4), pp. 216–220. doi:10.1177/089198879400700404.
- Li, C.T. et al. (2014). 'High dosage of hypnotics predicts subsequent sleep-related breathing disorders and is associated with worse outcomes for depression', *Sleep*, 37(4), pp. 803–809. doi:10.5665/sleep.3594.
- Li, W. et al. (2020). 'Association between sleep duration and quality and depressive symptoms among university students: A cross-sectional study', *PLoS ONE*, 15(9 September), pp. 1–10. doi:10.1371/journal.pone.0238811.
- Lim, J.A. et al. (2022). 'Greater variability in daily sleep efficiency predicts depression and anxiety in young adults: Estimation of depression severity using the two-week sleep quality records of wearable devices', *Frontiers in Psychiatry*, 13(November), pp. 1–10. doi:10.3389/fpsy.2022.1041747.
- Luis Torres , Manuela C Caciula , Alin S Tomoiaga, C.G.-G. (2023). 'Correlations between Mental Health, Physical Activity, and Body Composition in American College Students after the COVID-19 Pandemic Lockdown', *International Journal of Environmental Research and Public Health (IJERPH)*, 20(22), pp. 1–36. doi:10.3390/ijerph20227045.
- Luo, J. et al. (2024). 'Major Depressive Disorder Prediction Based on Sleep-Wake Disorders Symptoms in US Adolescents: A Machine Learning Approach from National Sleep Research Resource', *Psychology Research and Behavior Management*, 17(February), pp. 691–703. doi:10.2147/PRBM.S453046.
- Luo, J. & Lin, S. (2024). 'Sleep-wake changes and incident depressive symptoms in midlife women', *Scientific Reports*, 14(1), pp. 1–8. doi:10.1038/s41598-024-66145-3.
- Lv, X. et al. (2024). 'Body composition, lifestyle, and depression: a prospective study in the UK biobank', *BMC Public Health*, 24(1), pp. 1–10. doi:10.1186/s12889-024-17891-6.
- M Salih, A.M. et al. (2023). 'Exploring the relationship between insomnia and depression: a cross-sectional prospective study', *Annals of Medicine & Surgery*, 85(6), pp. 2573–2578. doi:10.1097/ms9.0000000000000784.
- Manuela C. Caciula, Luis Torres, S.A.T. (2024). 'A Comparative Study of Mental Health, Body Composition, and Physical Activity in American vs. Romanian College Students Post-COVID-19 Lockdown', *Journal of Exercise and Nutrition*, 7(1), pp. 1–8. doi:10.53520/jen2024.103171.
- Meitei, K.T. and Singh, H.S. (2019). 'Prevalence of Depression Among Indian Population-Citation Meitei KT, Singh HS. Prevalence of Depression Among Indian Population', *Online J Health Allied Scs*, 18(4), p. 12.



- Merino, M. et al. (2024). 'Body Perceptions and Psychological Well-Being: A Review of the Impact of Social Media and Physical Measurements on Self-Esteem and Mental Health with a Focus on Body Image Satisfaction and Its Relationship with Cultural and Gender Factors', *Healthcare*, 12(14), p. 1396. doi:10.3390/healthcare12141396.
- Mollayeva, T. et al. (2016). 'The Pittsburgh sleep quality index as a screening tool for sleep dysfunction in clinical and non-clinical samples: A systematic review and meta-analysis', *Sleep Medicine Reviews*, 25, pp. 52–73. doi:10.1016/j.smr.2015.01.009.
- Mu, F.Z. et al. (2024). 'Influence of physical exercise on negative emotions in college students: chain mediating role of sleep quality and self-rated health', *Frontiers in Public Health*, 12(May), pp. 1–14. doi:10.3389/fpubh.2024.1402801.
- Nakajima, S. et al. (2023). 'Transdiagnostic association between subjective insomnia and depressive symptoms in major psychiatric disorders', *Frontiers in Psychiatry*, 14(April), pp. 1–10. doi:10.3389/fpsy.2023.1114945.
- Nie, J., Zhang, W. & Liu, Y. (2017). 'Exploring depression, self-esteem and verbal fluency with different degrees of internet addiction among Chinese college students', *Comprehensive Psychiatry*, 72, pp. 114–120. doi:10.1016/j.comppsy.2016.10.006.
- O'Leary, D. et al. (2021). 'Subjective social status and physical health: The role of negative affect and reappraisal', *Social Science and Medicine*, 291(July), p. 114272. doi:10.1016/j.socscimed.2021.114272.
- Omichi, C. et al. (2022). 'Prolonged Sleep Latency and Reduced REM Latency Are Associated with Depressive Symptoms in a Japanese Working Population', *International Journal of Environmental Research and Public Health*, 19(4), pp. 1–10. doi:10.3390/ijerph19042112.
- Otte, J.L. & Carpenter, J.S. (2009). 'Theories, models, and frameworks related to sleep-wake disturbances in the context of cancer', *Cancer Nursing*, 32(2), pp. 90–104. doi:10.1097/01.NCC.0000339261.06442.7d.
- Riemann, D. et al. (2020). 'Sleep, insomnia, and depression', *Neuropsychopharmacology*, 45(1), pp. 74–89. doi:10.1038/s41386-019-0411-y.
- Rindler, G.A., Gries, A. and Freidl, W. (2023). 'Associations between overweight, obesity, and mental health: a retrospective study among European adults aged 50+', *Frontiers in Public Health*, 11(July). doi:10.3389/fpubh.2023.1206283.
- Salk, R.H., Hyde, J.S. & Abramson, L.Y. (2017). 'Gender differences in depression in representative national samples: Meta-analyses of diagnoses and symptoms', *Psychological Bulletin*, 143(8), pp. 783–822. doi:10.1037/bul0000102.
- Shahbaz Alam, Sheetal Kalra, Sadhu Charan Mohapatra, Richa Rai, D.R. 3 (2023). 'Frequency and Associated Factors of Physical Activity and Sleep with Depression in College Going Students', *Journal of Dow University of Health Sciences*, 17(1), pp. 17–24. doi:10.36570/jduhs.2023.1.1745.
- Shao, R. et al. (2020). 'Prevalence of depression and anxiety and correlations between depression, anxiety, family functioning, social support, and coping styles among Chinese medical students', *BMC Psychology*, 8(1), pp. 1–19. doi:10.1186/s40359-020-00402-8.



- Shim, E.J. et al. (2019). 'A longitudinal analysis of the relationships among daytime dysfunction, fatigue, and depression in college students', *Journal of American College Health*, 67(1), pp. 1–26. doi:10.1080/07448481.2018.1462819.
- Singh, B. et al. (2023). 'Effectiveness of physical activity interventions for improving depression, anxiety and distress: An overview of systematic reviews', *British Journal of Sports Medicine*, pp. 1203–1209. doi:10.1136/bjsports-2022-106195.
- Tanahira, C. et al. (2023). 'Bidirectional relationship between sleep and depression Shinnosuke', *Neuroscience Research*, 63(April), pp. 1–8. doi:10.1016/j.neures.2023.04.006.
- Vanteemar S. Sreeraj, S. et al. (2019). 'Indian nursing students' attitudes toward mental illness and persons with mental illness', *Industrial Psychiatry Journal*, 33(1), pp. 195–201. doi:10.4103/ipj.ipj.
- Viinikainen, J. et al. (2024). 'Effect of weight on depression using multiple genetic instruments', *PLoS ONE*, 19(2 FEBRUARY), pp. 1–8. doi:10.1371/journal.pone.0297594.
- William, W.K. (1965). 'Zung Self-Rating Depression Scale', *Encyclopedia of Quality of Life and Well-Being Research*, pp. 63–70. doi:10.1007/978-94-007-0753-5_104628.
- Wilson, K.T. et al. (2014). 'Social, behavioral, and sleep characteristics associated with depression symptoms among undergraduate students at a women's college: A cross-sectional depression survey, 2012', *BMC Women's Health*, 14(1). doi:10.1186/1472-6874-14-8.
- Wolf, M.R. and Rosenstock, J.B. (2017). 'Inadequate Sleep and Exercise Associated with Burnout and Depression among Medical Students', *Academic Psychiatry*, 41(2), pp. 174–179. doi:10.1007/s40596-016-0526-y.
- Xiang, M.Q. et al. (2020). 'Relationship of Physical Activity With Anxiety and Depression Symptoms in Chinese College Students During the COVID-19 Outbreak', *Frontiers in Psychology*, 11(November), pp. 1–7. doi:10.3389/fpsyg.2020.582436.
- Yan, B. et al. (2022). 'Sleep Efficiency May Predict Depression in a Large Population-Based Study', *Frontiers in Psychiatry*, 13(April), pp. 1–7. doi:10.3389/fpsyg.2022.838907.
- Yousef D Alqurashi, Ali H. Al Qattan, H.E.A.A. (2022). 'Association of sleep duration and quality with depression among university students and faculty'. Damman, Saudi Arabia: *Acta Biomed*, pp. 1–8. doi:10.23750/abm.v93i5.13002.
- Yuan, W. et al. (2023). 'Association of body fat distribution with depression and social anxiety in children and adolescents: A cross-sectional study based on dual-energy X-ray detection', *Beijing da xue xue bao. Yi xue ban = Journal of Peking University. Health sciences*, 55(3), pp. 429–435. doi:10.19723/j.issn.1671-167X.2023.03.007.
- Zhang, D. et al. (2023). 'Association between healthy sleep patterns and depressive trajectories among college students: a prospective cohort study', *BMC Psychiatry*, 23(1), pp. 1–10. doi:10.1186/s12888-023-04596-0.
- Zhang, J., Zheng, S. & Hu, Z. (2022). 'The Effect of Physical Exercise on Depression in College Students: The Chain Mediating Role of Self-Concept and Social Support', *Frontiers in Psychology*, 13(May), pp. 1–6. doi:10.3389/fpsyg.2022.841160.



Zhang, R., Cheng, M. & Zhao, L. (2024). 'An investigative study of anxiety, depression, and sleep quality in college students after the COVID-19 pandemic - the moderating effect of physical activity', *Journal of Medicine and Health Sciences Research*, 7(3), pp. 10–15. doi:10.21839/jmhsr.2024.v7.8779.

Zhou, H. et al. (2021). 'Association of sedentary behavior and physical activity with depression in sport university students', *International Journal of Environmental Research and Public Health*, 18(18). doi:10.3390/ijerph18189881.